

Summary of the Ontario Publicly Funded Immunization Schedule for Children Starting Immunization in Infancy

Date issued: July 2020

NOTE: CHILDREN ARE REQUIRED TO BE UP TO DATE FOR THEIR AGE TO ATTEND CHILD CARE AND SCHOOL

Age	Vaccines to be given	Route for Administration
2 months	DTaP-IPV-HIB (Pediacel) Pneumococcal Conjugate 13 (Prevnar-13) Rot-5 (RotaTeq)	IM – vastus lateralis (leg) IM – vastus lateralis (leg) Oral
4 months	DTaP-IPV-HIB (Pediacel) Pneumococcal Conjugate 13 (Prevnar-13) Rot-5 (RotaTeq)	IM – vastus lateralis (leg) IM – vastus lateralis (leg) Oral
6 months	DTaP-IPV-HIB (Pediacel) Rot-5 (RotaTeq)	IM – vastus lateralis (leg) Oral
12 months	MMR (MMRII, Priorix)* Meningococcal Conjugate C (Neis-Vac-C, Menjugate)** Pneumococcal Conjugate 13 (Prevnar-13)	SC – upper outer tricep area IM – deltoid IM - deltoid
15 months	Varicella (Varivax III, Varilrix)*	SC – upper outer tricep area
18 months	DTaP-IPV-HIB (Pediacel)	IM - deltoid
4-6 years	MMRV (Priorix-Tetra, ProQuad)* Tdap-IPV (Adacel-Polio, Boostrix-Polio)***	SC – upper outer tricep area IM - deltoid
11-12 years (given in school in Grade 7)	Hepatitis B (Enerix, Recombivax) (1.0 ml dose) HPV-9 (Gardasil 9) Meningococcal Conjugate ACYW-135 (Menactra)	IM - deltoid (2 doses, 6 months apart) IM - deltoid (2 doses, 6 months apart) IM - deltoid
10 years after Tdap-IPV	Tdap (Adacel, Boostrix)	IM – deltoid

* All live vaccines must be given **on or after** the 1st birthday (MMR, Varicella, MMRV). The minimum spacing between live vaccines is 4 weeks except when BOTH vaccines contain varicella, when the minimum spacing is 6 weeks.

** Men-C-C vaccine must be given **on or after** the 1st birthday.

*** Tdap-IPV (5th dose of these antigens) must be given **on or after** the 4th birthday.

CONSULT THE FULL PUBLICLY FUNDED IMMUNIZATION SCHEDULES FOR ONTARIO FOR MORE INFORMATION ON DOSING, SPACING, CATCH UP SERIES AND OTHER VACCINE REQUIREMENTS.

THE PUBLICLY FUNDED SCHEDULE IS SUBJECT TO CHANGE AT ANY TIME.