

References

1. David Hammond et. al. Prevalence of vaping and smoking among adolescents in Canada, England, and the United States: repeat national cross-sectional surveys. | BMJ 2019;365:l2219 | doi: 10.1136/bmj.l2219.
2. Health Canada. Canadian student tobacco, alcohol and drugs (CSTADS) survey 2018-2019. aem. Published December 19, 2019. <https://www.canada.ca/en/health-canada/services/canadian-student-tobacco-alcohol-drugs-survey/2018-2019-detailed-tables.html>.
3. Yuan M, Cross SJ, Loughlin SE, Leslie FM. Nicotine and the adolescent brain: Nicotine and the adolescent brain. J Physiol. 2015;593(16):3397-3412. doi:10.1113/JP2704922328.
4. Heart & Stroke Funded Survey, The 2020-2021 Youth and Young Adults Vaping Project, 2021.
5. CDC Foundation, Data Brief March 2021: Monitoring U.S. E-Cigarette Sales: National Trends retrieved from <https://www.cdcfoundation.org/National-E-CigaretteSales-DataBrief-2021-Mar21?inline>.
6. Berry, K. M., Fetterman, J. L., Benjamin, E. J, et al. (2019). Association of Electronic Cigarette Use With Subsequent Initiation of Tobacco Cigarettes in US Youths. JAMA Netw Open, 20192(2):e187794. doi:10.1001/jamanetworkopen.2018.7794. Retrieved from <https://jamanetwork.com/journals/jamanetworkopen/fullarticle/2723425>.
7. Osibogun O, Bursac Z, Maziak W. E-Cigarette Use and Regular Cigarette Smoking Among Youth: Population Assessment of Tobacco and Health Study (2013-2016). *Am J Prev Med*. 2020;58(5):657-665. doi:10.1016/j.amepre.2020.01.003.
8. Zare S, Nemati M, Zheng Y. A systematic review of consumer preference for e-cigarette attributes: Flavor, nicotine strength, and type. *PLOS ONE*. 2018;13:e0194145. doi:10.1371/journal.pone.0194145.